

Chapter 9: Physiotherapy Modalities: Ancillary and Preparatory to the Chiropractic Adjustment

Introduction

Physical modalities utilized in healthcare can be traced back from Hippocrates around 400BC, and Galen, who was considered the “Father of Sports Medicine”, as he treated the Roman gladiators circa 200AD (6). Since, the dichotomy within the healing arts between physical modalities of treatment and chemical modalities has taken turns in dominance throughout history.

The introduction of physiotherapy within the United States can be best associated with two phenomena within American history. The first of these events being the epidemics of poliomyelitis during the 1800’s through the 1950’s, and the onset of World War I and World War II (4). These two events left patients with unique symptoms and impairments that responded in a superior manor to physical modalities such as exercise, hydrotherapy and massage; rather than the conventional medical treatments at that time.

Chiropractic Physiotherapy is defined as such:

“Chiropractic physiologic therapeutics encompasses the diagnosis and treatment of disorders of the body using the natural forces of healing, such as air, cold, electricity, rest, exercise, traction, heat, light, massage, water and other forces of nature.”(This is the ACA definition so may want to replace with an ICA or more neutral one)(2).

The use of physical modalities within chiropractic, as well as within the physical therapy profession, has occurred concomitantly. Within chiropractic, physiotherapy modalities can be seen early in the profession’s history. While some postulate that physiotherapy in chiropractic practice can be traced back to D.D. Palmer’s magnetic healing practice which incorporated a sort of “massage”(5). A more deliberate union was made in 1914 when National Chiropractic College incorporated physiotherapy in its program(1). Photos of the B.J. Palmer Chiropractic Research Clinic, circa 1945, show a state of the art rehabilitation lab containing the latest physiotherapy equipment of the time(5). Since, chiropractic has enjoyed the benefit of including physiotherapy into their scope of practice.

The types of modalities utilized in physiotherapy have been scrutinized for their lack of “evidence” in the past. As a result the American Physical Therapy Association (APTA) has created “Hooked on Evidence”. This program, in association with “Vision 2020”, and 187 Doctor of Physical Therapy (DPT) programs countrywide are an attempt by the physical therapy profession to gain autonomy and direct access privileges in the ever-growing market for physical medicine utilization(3). It is critical that the chiropractic profession stakes its claim to this area of professional overlap in a way that that is defined and substantiated.

In the new era of “evidence-based” chiropractic practice, it is imperative that the profession remain vigilant in the appraisal of the scientific literature as it pertains to chiropractic practice. This is especially crucial when ascribing validity to the various modalities.

Several decades ago, the Board of Directors of the International Chiropractors Association (ICA) voted to include a variety of physiotherapy modalities as used by chiropractors. These procedures were accepted if their use was “ancillary or preparatory” to the chiropractic spinal adjustment. While the major method of chiropractic care, recognized by the ICA, is the chiropractic spinal adjustment, the ICA realizes that certain modalities, when used in conjunction with the adjustment, may enhance healing and reduce the suffering of chiropractic patients.

While it is difficult to determine the current evidence for each modality used by chiropractors, several of the most common physiotherapy modalities were searched within the Medline Database. As discussed in previous chapters, there are four major levels of evidence recognized by the United States Department of Health and Human Services:

- **Level 1.** *Randomized controlled/clinical trials*—includes quasi-randomized processes such as alternate allocation.
- **Level 2.** *Non-randomized controlled/clinical trial*—a prospective (pre-planned) study, with predetermined eligibility criteria and outcome measures.
- **Level 3.** *Observational studies with controls*—includes retrospective, interrupted time series (a change in trend attributable to the intervention), case-control studies, cohort studies with controls, and health services research that includes adjustment for likely confounding variables.
- **Level 4.** *Observational studies without controls* (e.g., cohort studies without controls, case series without controls, and case studies without controls).

Before discussing the Levels of Evidence for each common modality, Table 1 is provided as a summary of the evidence for each one.

Table 1

	Modalities	Level I	Level II	Level III	Level IV	Total Number of References	Rating
22.	Cold/Cryotherapy +	4	0	0	2	6	A
23.	Cold/Cryotherapy -	2	1	0	0	3	A
24.	Superficial Heat +	7	0	0	2	9	B
25.	Superficial Heat -	1	0	0	0	1	A
26.	Therapeutic Exercise +	71	10	6	44	131	A
27.	Therapeutic Exercise -	3	0	0	1	4	A
28.	Low Level Laser +	27	1	1	7	36	A
29.	Low Level Laser -	13	0	0	0	13	A
30.	TENS +	41	1	1	34	77	A
31.	TENS -	12	0	0	1	13	A
32.	Whole Body Vibration +	10	0	2	1	13	B
33.	Whole Body Vibration -	1	0	0	0	1	A
34.	Yoga +	19	1	2	9	31	A
35.	Yoga -	2	0	0	0	2	A
36.	Extension Traction +	1	3	0	3	7	A
37.	Extension Traction -	0	0	0	0	0	A

Discussion:

Cold/Cryotherapy:

Positive:

Level 1: References:

C. E. Gibbons, M. C. Solan, D. M. Ricketts, and M. Patterson. Cryotherapy compared with Robert Jones bandage after total knee replacement: a prospective randomized trial. *Int.Orthop.* 25 (4):250-252, 2001.

J. Hochberg. A randomized prospective study to assess the efficacy of two cold-therapy treatments following carpal tunnel release. *J.Hand Ther.* 14 (3):208-215, 2001.

A. Holmstrom and B. C. Hardin. Cryo/Cuff compared to epidural anesthesia after knee unicompartmental arthroplasty: a prospective, randomized and controlled study of 60 patients with a 6-week follow-up. *J.Arthroplasty* 20 (3):316-321, 2005.

O. K. Jensen, F. F. Nielsen, and L. Vosmar. An open study comparing manual therapy with the use of cold packs in the treatment of post-traumatic headache. *Cephalalgia* 10 (5):241-250, 1990.

Level 2-3: (0 references)

Level 4:

C. R. Green, A. M. de Rosayro, and A. R. Tait. The role of cryoanalgesia for chronic thoracic pain: results of a long-term follow up. *J.Natl.Med.Assoc.* 94 (8):716-720, 2002.

E. Morsi. Continuous-flow cold therapy after total knee arthroplasty. *J.Arthroplasty* 17 (6):718-722, 2002.

Cold/Cryotherapy:

Negative:

Level 1:

W. L. Healy, J. Seidman, B. A. Pfeifer, and D. G. Brown. Cold compressive dressing after total knee arthroplasty. *Clin.Orthop.Relat Res.* (299):143-146, 1994.

H. E. Hirvonen, M. K. Mikkelsen, H. Kautiainen, T. H. Pohjolainen, and M. Leirisalo-Repo. Effectiveness of different cryotherapies on pain and disease activity in active rheumatoid arthritis. A randomised single blinded controlled trial. *Clin.Exp.Rheumatol.* 24 (3):295-301, 2006.

Level 2:

B. Kullenberg, S. Ylipaa, K. Soderlund, and S. Resch. Postoperative cryotherapy after total knee arthroplasty: a prospective study of 86 patients. *J.Arthroplasty* 21 (8):1175-1179, 2006.

Superficial Heat:

Positive:

Level 1:

B. Kettenmann, C. Wille, E. Lurie-Luke, D. Walter, and G. Kobal. Impact of continuous low level heatwrap therapy in acute low back pain patients: subjective and objective measurements. *Clin.J.Pain* 23 (8):663-668, 2007.

J. M. Mayer, L. Ralph, M. Look, G. N. Erasala, J. L. Verna, L. N. Matheson, and V. Mooney. Treating acute low back pain with continuous low-level heat wrap therapy and/or exercise: a randomized controlled trial. *Spine J.* 5 (4):395-403, 2005.

J. M. Mayer, V. Mooney, L. N. Matheson, G. N. Erasala, J. L. Verna, B. E. Udermann, and S. Leggett. Continuous low-level heat wrap therapy for the prevention and early phase treatment of delayed-onset muscle soreness of the low back: a randomized controlled trial. *Arch.Phys.Med.Rehabil.* 87 (10):1310-1317, 2006.

S. F. Nadler, D. J. Steiner, G. N. Erasala, D. A. Hengehold, R. T. Hinkle, Goodale M. Beth, S. B. Abeln, and K. W. Weingand. Continuous low-level heat wrap therapy provides more efficacy than Ibuprofen and acetaminophen for acute low back pain. *Spine* 27 (10):1012-1017, 2002.

S. F. Nadler, D. J. Steiner, G. N. Erasala, D. A. Hengehold, S. B. Abeln, and K. W. Weingand. Continuous low-level heatwrap therapy for treating acute nonspecific low back pain. *Arch.Phys.Med.Rehabil.* 84 (3):329-334, 2003.

S. F. Nadler, D. J. Steiner, S. R. Petty, G. N. Erasala, D. A. Hengehold, and K. W. Weingand. Overnight use of continuous low-level heatwrap therapy for relief of low back pain. *Arch.Phys.Med.Rehabil.* 84 (3):335-342, 2003.

X. G. Tao and E. J. Bernacki. A randomized clinical trial of continuous low-level heat therapy for acute muscular low back pain in the workplace. *J.Occup.Environ.Med.* 47 (12):1298-1306, 2005.

Level 2-3: (0 references)

Level 4:

B. Curkovic, V. Vitulic, D. Babic-Naglic, and T. Durrigl. The influence of heat and cold on the pain threshold in rheumatoid arthritis. *Z.Rheumatol.* 52 (5):289-291, 1993.

A. Lloyd, D. A. Scott, R. L. Akehurst, E. Lurie-Luke, and G. Jessen. Cost-effectiveness of low-level heat wrap therapy for low back pain. *Value.Health* 7 (4):413-422, 2004.

Superficial Heat:

Negative:

Level 1:

Y. Saeki. Effect of local application of cold or heat for relief of pricking pain. *Nurs.Health Sci.* 4 (3):97-105, 2002.

Level 2-4: (0 references)

Therapeutic Exercise:

Positive:

Level 1:

F. Avraham, S. Aviv, P. Ya'akobi, H. Faran, Z. Fisher, Y. Goldman, G. Neeman, and E. Carmeli. The efficacy of treatment of different intervention programs for patellofemoral pain syndrome--a single blinded randomized clinical trial. Pilot study. *ScientificWorldJournal.* 7:1256-1262, 2007.

M. D. Bang and G. D. Deyle. Comparison of supervised exercise with and without manual physical therapy for patients with shoulder impingement syndrome. *J.Orthop.Sports Phys.Ther.* 30 (3):126-137, 2000.

H. Bentsen, F. Lindgarde, and R. Manthorpe. The effect of dynamic strength back exercise and/or a home training program in 57-year-old women with chronic low back pain. Results of a prospective randomized study with a 3-year follow-up period. *Spine* 22 (13):1494-1500, 1997.

O. H. Bjarnadottir, A. D. Konradsdottir, K. Reynisdottir, and E. Olafsson. Multiple sclerosis and brief moderate exercise. A randomised study. *Mult.Scler.* 13 (6):776-782, 2007.

L. Bunketorp, M. Lindh, J. Carlsson, and E. Stener-Victorin. The effectiveness of a supervised physical training model tailored to the individual needs of patients with whiplash-associated disorders--a randomized controlled trial. *Clin.Rehabil.* 20 (3):201-217, 2006.

T. T. Chiu, T. H. Lam, and A. J. Hedley. A randomized controlled trial on the efficacy of exercise for patients with chronic neck pain. *Spine* 30 (1):E1-E7, 2005.

T. Cochrane, R. C. Davey, and S. M. Matthes Edwards. Randomised controlled trial of the cost-effectiveness of water-based therapy for lower limb osteoarthritis. *Health Technol.Assess.* 9 (31):iii-xi, 1, 2005.

G. E. Djavid, R. Mehrdad, M. Ghasemi, H. Hasan-Zadeh, A. Sotoodeh-Manesh, and G. Pouryaghoub. In chronic low back pain, low level laser therapy combined with exercise is more beneficial than exercise alone in the long term: a randomised trial. *Aust.J.Physiother.* 53 (3):155-160, 2007.

K. J. Dodd, N. F. Taylor, and H. K. Graham. A randomized clinical trial of strength training in young people with cerebral palsy. *Dev.Med.Child Neurol.* 45 (10):652-657, 2003.

V. E. Drory, E. Goltsman, J. G. Reznik, A. Mosek, and A. D. Korczyn. The value of muscle exercise in patients with amyotrophic lateral sclerosis. *J.Neurol.Sci.* 191 (1-2):133-137, 2001.

P. Duncan, L. Richards, D. Wallace, J. Stoker-Yates, P. Pohl, C. Luchies, A. Ogle, and S. Studenski. A randomized, controlled pilot study of a home-based exercise program for individuals with mild and moderate stroke. *Stroke* 29 (10):2055-2060, 1998.

K. Dzedzic, J. Hill, M. Lewis, J. Sim, J. Daniels, and E. M. Hay. Effectiveness of manual therapy or pulsed shortwave diathermy in addition to advice and exercise for neck disorders: a pragmatic randomized controlled trial in physical therapy clinics. *Arthritis Rheum.* 53 (2):214-222, 2005.

H. Epps, L. Ginnelly, M. Utley, T. Southwood, S. Gallivan, M. Sculpher, and P. Woo. Is hydrotherapy cost-effective? A randomised controlled trial of combined hydrotherapy programmes compared with physiotherapy land techniques in children with juvenile idiopathic arthritis. *Health Technol.Assess.* 9 (39):iii-x, 1, 2005.

L. Eversden, F. Maggs, P. Nightingale, and P. Jobanputra. A pragmatic randomised controlled trial of hydrotherapy and land exercises on overall well being and quality of life in rheumatoid arthritis. *BMC.Musculoskelet.Disord.* 8:23, 2007.

D. Falla, G. Jull, P. Hodges, and B. Vicenzino. An endurance-strength training regime is effective in reducing myoelectric manifestations of cervical flexor muscle fatigue in females with chronic neck pain. *Clin.Neurophysiol.* 117 (4):828-837, 2006.

C. Fernandez-de-Las-Penas, C. Alonso-Blanco, M. Morales-Cabezas, and J. C. Miangolarra-Page. Two exercise interventions for the management of patients with ankylosing spondylitis: a randomized controlled trial. *Am.J.Phys.Med.Rehabil.* 84 (6):407-419, 2005.

C. Fernandez-de-Las-Penas, C. Alonso-Blanco, I. M. Alguacil-Diego, and J. C. Miangolarra-Page. One-year follow-up of two exercise interventions for the management of patients with ankylosing spondylitis: a randomized controlled trial. *Am.J.Phys.Med.Rehabil.* 85 (7):559-567, 2006.

W. R. Ferrell, N. Tennant, R. D. Sturrock, L. Ashton, G. Creed, G. Brydson, and D. Rafferty. Amelioration of symptoms by enhancement of proprioception in patients with joint hypermobility syndrome. *Arthritis Rheum.* 50 (10):3323-3328, 2004.

G. K. Fitzgerald, M. J. Axe, and L. Snyder-Mackler. The efficacy of perturbation training in nonoperative anterior cruciate ligament rehabilitation programs for physical active individuals. *Phys.Ther.* 80 (2):128-140, 2000.

N. E. Foster, E. Thomas, P. Barlas, J. C. Hill, J. Young, E. Mason, and E. M. Hay. Acupuncture as an adjunct to exercise based physiotherapy for osteoarthritis of the knee: randomised controlled trial. *BMJ* 335 (7617):436, 2007.

K. Y. Fulcher and P. D. White. Randomised controlled trial of graded exercise in patients with the chronic fatigue syndrome. *BMJ* 314 (7095):1647-1652, 1997.

L. J. Goldby, A. P. Moore, J. Doust, and M. E. Trew. A randomized controlled trial investigating the efficiency of musculoskeletal physiotherapy on chronic low back disorder. *Spine* 31 (10):1083-1093, 2006.

F. Guler-Uysal and E. Kozanoglu. Comparison of the early response to two methods of rehabilitation in adhesive capsulitis. *Swiss.Med.Wkly.* 134 (23-24):353-358, 2004.

R. S. Hinman, S. E. Heywood, and A. R. Day. Aquatic physical therapy for hip and knee osteoarthritis: results of a single-blind randomized controlled trial. *Phys.Ther.* 87 (1):32-43, 2007.

M. A. Hirsch, T. Toole, C. G. Maitland, and R. A. Rider. The effects of balance training and high-intensity resistance training on persons with idiopathic Parkinson's disease. *Arch.Phys.Med.Rehabil.* 84 (8):1109-1117, 2003.

H. L. Hoeksma, J. Dekker, H. K. Ronday, A. Heering, Lubbe N. van der, C. Vel, F. C. Breedveld, and C. H. van den Ende. Comparison of manual therapy and exercise therapy in osteoarthritis of the hip: a randomized clinical trial. *Arthritis Rheum.* 51 (5):722-729, 2004.

A. Jordan, T. Bendix, H. Nielsen, F. R. Hansen, D. Host, and A. Winkel. Intensive training, physiotherapy, or manipulation for patients with chronic neck pain. A prospective, single-blinded, randomized clinical trial. *Spine* 23 (3):311-318, 1998.

G. Jull, P. Trott, H. Potter, G. Zito, K. Niere, D. Shirley, J. Emberson, I. Marschner, and C. Richardson. A randomized controlled trial of exercise and manipulative therapy for cervicogenic headache. *Spine* 27 (17):1835-1843, 2002.

G. Jull, D. Falla, J. Treleaven, P. Hodges, and B. Vicenzino. Retraining cervical joint position sense: the effect of two exercise regimes. *J.Orthop.Res.* 25 (3):404-412, 2007.

M. Kankaanpaa, S. Taimela, O. Airaksinen, and O. Hanninen. The efficacy of active rehabilitation in chronic low back pain. Effect on pain intensity, self-experienced disability, and lumbar fatigability. *Spine* 24 (10):1034-1042, 1999.

D. J. Kidgell, D. M. Horvath, B. M. Jackson, and P. J. Seymour. Effect of six weeks of dura disc and mini-trampoline balance training on postural sway in athletes with functional ankle instability. *J.Strength.Cond.Res.* 21 (2):466-469, 2007.

S. H. Kim, K. I. Ha, M. W. Jung, M. S. Lim, Y. M. Kim, and J. H. Park. Accelerated rehabilitation after arthroscopic Bankart repair for selected cases: a prospective randomized clinical study. *Arthroscopy* 19 (7):722-731, 2003.

J. S. Lewis, J. S. Hewitt, L. Billington, S. Cole, J. Byng, and S. Karayiannis. A randomized clinical trial comparing two physiotherapy interventions for chronic low back pain. *Spine* 30 (7):711-721, 2005.

M. Lewis, M. James, E. Stokes, J. Hill, J. Sim, E. Hay, and K. Dziedzic. An economic evaluation of three physiotherapy treatments for non-specific neck disorders alongside a randomized trial. *Rheumatology.(Oxford)* 46 (11):1701-1708, 2007.

A. E. Ljunggren, H. Weber, O. Kogstad, E. Thom, and G. Kirkesola. Effect of exercise on sick leave due to low back pain. A randomized, comparative, long-term study. *Spine* 22 (14):1610-1616, 1997.

P. M. Ludewig and J. D. Borstad. Effects of a home exercise programme on shoulder pain and functional status in construction workers. *Occup. Environ. Med.* 60 (11):841-849, 2003.

- B. Malmros, L. Mortensen, M. B. Jensen, and P. Charles. Positive effects of physiotherapy on chronic pain and performance in osteoporosis. *Osteoporos.Int.* 8 (3):215-221, 1998.
- T. Manini, M. Marko, T. VanArnam, S. Cook, B. Fernhall, J. Burke, and L. Ploutz-Snyder. Efficacy of resistance and task-specific exercise in older adults who modify tasks of everyday life. *J.Gerontol.A Biol.Sci.Med.Sci.* 62 (6):616-623, 2007.
- A. F. Mannion, M. Muntener, S. Taimela, and J. Dvorak. A randomized clinical trial of three active therapies for chronic low back pain. *Spine* 24 (23):2435-2448, 1999.
- A. F. Mannion, M. Muntener, S. Taimela, and J. Dvorak. Comparison of three active therapies for chronic low back pain: results of a randomized clinical trial with one-year follow-up. *Rheumatology.(Oxford)* 40 (7):772-778, 2001.
- L. Martin, A. Nutting, B. R. MacIntosh, S. M. Edworthy, D. Butterwick, and J. Cook. An exercise program in the treatment of fibromyalgia. *J.Rheumatol.* 23 (6):1050-1053, 1996.
- C. J. McCarthy, P. M. Mills, R. Pullen, G. Richardson, N. Hawkins, C. R. Roberts, A. J. Silman, and J. A. Oldham. Supplementation of a home-based exercise programme with a class-based programme for people with osteoarthritis of the knees: a randomised controlled trial and health economic analysis. *Health Technol.Assess.* 8 (46):iii-61, 2004.
- M. Mulet, K. L. Decker, J. O. Look, P. A. Lenton, and E. L. Schiffman. A randomized clinical trial assessing the efficacy of adding 6 x 6 exercises to self-care for the treatment of masticatory myofascial pain. *J.Orofac.Pain* 21 (4):318-328, 2007.
- M. E. Nelson, M. A. Fiatarone, C. M. Morganti, I. Trice, R. A. Greenberg, and W. J. Evans. Effects of high-intensity strength training on multiple risk factors for osteoporotic fractures. A randomized controlled trial. *JAMA* 272 (24):1909-1914, 1994.
- L. Niemisto, P. Rissanen, S. Sarna, T. Lahtinen-Suopanki, K. A. Lindgren, and H. Hurri. Cost-effectiveness of combined manipulation, stabilizing exercises, and physician consultation compared to physician consultation alone for chronic low back pain: a prospective randomized trial with 2-year follow-up. *Spine* 30 (10):1109-1115, 2005.
- S. O'Leary, D. Falla, P. W. Hodges, G. Jull, and B. Vicenzino. Specific therapeutic exercise of the neck induces immediate local hypoalgesia. *J.Pain* 8 (11):832-839, 2007.
- S. D. O'Shea, N. F. Taylor, and J. D. Paratz. A predominantly home-based progressive resistance exercise program increases knee extensor strength in the short-term in people with chronic obstructive pulmonary disease: a randomised controlled trial. *Aust.J.Physiother.* 53 (4):229-237, 2007.
- L. H. Pengel, K. M. Refshauge, C. G. Maher, M. K. Nicholas, R. D. Herbert, and P. McNair. Physiotherapist-directed exercise, advice, or both for subacute low back pain: a randomized trial. *Ann.Intern.Med.* 146 (11):787-796, 2007.
- T. Petersen, P. Kryger, C. Ekdahl, S. Olsen, and S. Jacobsen. The effect of McKenzie therapy as compared with that of intensive strengthening training for the treatment of patients with subacute or chronic low back pain: A randomized controlled trial. *Spine* 27 (16):1702-1709, 2002.

S. H. Peurala, I. M. Tarkka, K. Pitkanen, and J. Sivenius. The effectiveness of body weight-supported gait training and floor walking in patients with chronic stroke. *Arch.Phys.Med.Rehabil.* 86 (8):1557-1564, 2005.

R. M. Rine, J. Braswell, D. Fisher, K. Joyce, K. Kalar, and M. Shaffer. Improvement of motor development and postural control following intervention in children with sensorineural hearing loss and vestibular impairment. *Int.J.Pediatr.Otorhinolaryngol.* 68 (9):1141-1148, 2004.

H. Rogind, B. Bibow-Nielsen, B. Jensen, H. C. Moller, H. Frimodt-Moller, and H. Bliddal. The effects of a physical training program on patients with osteoarthritis of the knees. *Arch.Phys.Med.Rehabil.* 79 (11):1421-1427, 1998.

G. Senbursa, G. Baltaci, and A. Atay. Comparison of conservative treatment with and without manual physical therapy for patients with shoulder impingement syndrome: a prospective, randomized clinical trial. *Knee.Surg.Sports Traumatol.Arthrosc.* 15 (7):915-921, 2007.

C. Sherrington, S. R. Lord, and R. D. Herbert. A randomised trial of weight-bearing versus non-weight-bearing exercise for improving physical ability in inpatients after hip fracture. *Aust.J.Physiother.* 49 (1):15-22, 2003.

T. Sjogren, K. J. Nissinen, S. K. Jarvenpaa, M. T. Ojanen, H. Vanharanta, and E. A. Malkia. Effects of a workplace physical exercise intervention on the intensity of headache and neck and shoulder symptoms and upper extremity muscular strength of office workers: a cluster randomized controlled cross-over trial. *Pain* 116 (1-2):119-128, 2005.

R. Song, E. O. Lee, P. Lam, and S. C. Bae. Effects of tai chi exercise on pain, balance, muscle strength, and perceived difficulties in physical functioning in older women with osteoarthritis: a randomized clinical trial. *J.Rheumatol.* 30 (9):2039-2044, 2003.

L. M. Spencer, J. A. Alison, and Z. J. McKeough. Do supervised weekly exercise programs maintain functional exercise capacity and quality of life, twelve months after pulmonary rehabilitation in COPD? *BMC.Pulm.Med.* 7:7, 2007.

W. R. Stanton and G. A. Jull. Cervicogenic headache: locus of control and success of treatment. *Headache* 43 (9):956-961, 2003.

D. Stasinopoulos and I. Stasinopoulos. Comparison of effects of exercise programme, pulsed ultrasound and transverse friction in the treatment of chronic patellar tendinopathy. *Clin.Rehabil.* 18 (4):347-352, 2004.

M. J. Stewart, C. G. Maher, K. M. Refshauge, R. D. Herbert, N. Bogduk, and M. Nicholas. Randomized controlled trial of exercise for chronic whiplash-associated disorders. *Pain* 128 (1-2):59-68, 2007.

K. Storheim, I. Holm, R. Gunderson, J. I. Brox, and K. Bo. The effect of comprehensive group training on cross-sectional area, density, and strength of paraspinal muscles in patients sick-listed for subacute low back pain. *J.Spinal Disord.Tech.* 16 (3):271-279, 2003.

A. Suputtitada, P. Yooktanon, and T. Rarerng-Ying. Effect of partial body weight support treadmill training in chronic stroke patients. *J.Med.Assoc.Thai.* 87 Suppl 2:S107-S111, 2004.

S. Taimela, E. P. Takala, T. Asklof, K. Seppala, and S. Parviainen. Active treatment of chronic neck pain: a prospective randomized intervention. *Spine* 25 (8):1021-1027, 2000.

T. A. Torstensen, A. E. Ljunggren, H. D. Meen, E. Odland, P. Mowinckel, and S. Geijerstam. Efficiency and costs of medical exercise therapy, conventional physiotherapy, and self-exercise in patients with chronic low back pain. A pragmatic, randomized, single-blinded, controlled trial with 1-year follow-up. *Spine* 23 (23):2616-2624, 1998.

T. Tritilanunt and W. Wajanavisit. The efficacy of an aerobic exercise and health education program for treatment of chronic low back pain. *J.Med.Assoc.Thai.* 84 Suppl 2:S528-S533, 2001.

H. van Ettekovon and C. Lucas. Efficacy of physiotherapy including a craniocervical training programme for tension-type headache; a randomized clinical trial. *Cephalalgia* 26 (8):983-991, 2006.

B. Wolf, H. Feys, Weerdt De, Meer J. van der, M. Noom, G. Aufdemkampe, and M. Noom. Effect of a physical therapeutic intervention for balance problems in the elderly: a single-blind, randomized, controlled multicentre trial. *Clin.Rehabil.* 15 (6):624-636, 2001.

J. Ylinen, E. P. Takala, M. Nykanen, A. Hakkinen, E. Malkia, T. Pohjolainen, S. L. Karppi, H. Kautiainen, and O. Airaksinen. Active neck muscle training in the treatment of chronic neck pain in women: a randomized controlled trial. *JAMA* 289 (19):2509-2516, 2003.

J. Ylinen, A. Hakkinen, M. Nykanen, H. Kautiainen, and E. P. Takala. Neck muscle training in the treatment of chronic neck pain: a three-year follow-up study. *Eura.Medicophys.* 43 (2):161-169, 2007.

J. J. Ylinen, A. H. Hakkinen, E. P. Takala, M. J. Nykanen, H. J. Kautiainen, E. A. Malkia, T. H. Pohjolainen, S. L. Karppi, and O. V. Airaksinen. Effects of neck muscle training in women with chronic neck pain: one-year follow-up study. *J.Strength.Cond.Res.* 20 (1):6-13, 2006.

J. J. Ylinen, E. P. Takala, M. J. Nykanen, H. J. Kautiainen, A. H. Hakkinen, and O. V. Airaksinen. Effects of twelve-month strength training subsequent to twelve-month stretching exercise in treatment of chronic neck pain. *J.Strength.Cond.Res.* 20 (2):304-308, 2006.

Level 2:

D. Borenstein. Prevalence and treatment outcome of primary and secondary fibromyalgia in patients with spinal pain. *Spine* 20 (7):796-800, 1995.

M. Buchner, A. Zahlten-Hinguranage, M. Schiltenswolf, and E. Neubauer. Therapy outcome after multidisciplinary treatment for chronic neck and chronic low back pain: a prospective clinical study in 365 patients. *Scand.J.Rheumatol.* 35 (5):363-367, 2006.

K. Chan, L. Qin, M. Lau, J. Woo, S. Au, W. Choy, K. Lee, and S. Lee. A randomized, prospective study of the effects of Tai Chi Chun exercise on bone mineral density in postmenopausal women. *Arch.Phys.Med.Rehabil.* 85 (5):717-722, 2004.

L. M. Giangregorio, A. L. Hicks, C. E. Webber, S. M. Phillips, B. C. Craven, J. M. Bugaresti, and N. McCartney. Body weight supported treadmill training in acute spinal cord injury: impact on muscle and bone. *Spinal Cord.* 43 (11):649-657, 2005.

K. R. Gottshall, M. E. Hoffer, R. J. Moore, and B. J. Balough. The role of vestibular rehabilitation in the treatment of Meniere's disease. *Otolaryngol.Head Neck Surg.* 133 (3):326-328, 2005.

D. E. Harrison, R. Cailliet, J. Betz, J. W. Haas, D. D. Harrison, T. J. Janik, and B. Holland. Conservative methods for reducing lateral translation postures of the head: a nonrandomized clinical control trial. *J.Rehabil.Res.Dev.* 41 (4):631-639, 2004.

A. Helewa, C. H. Goldsmith, H. A. Smythe, P. Lee, K. Obright, and L. Stitt. Effect of therapeutic exercise and sleeping neck support on patients with chronic neck pain: a randomized clinical trial. *J.Rheumatol.* 34 (1):151-158, 2007.

H. Karapolat, Y. Akkoc, I. Sari, S. Eyigor, S. Akar, Y. Kirazli, and N. Akkoc. Comparison of group-based exercise versus home-based exercise in patients with ankylosing spondylitis: effects on Bath Ankylosing Spondylitis Indices, quality of life and depression. *Clin.Rheumatol.*, 2007.

A. C. Kronhed and M. Moller. Effects of physical exercise on bone mass, balance skill and aerobic capacity in women and men with low bone mineral density, after one year of training--a prospective study. *Scand.J.Med.Sci.Sports* 8 (5 Pt 1):290-298, 1998.

J. K. Richardson, D. Sandman, and S. Vela. A focused exercise regimen improves clinical measures of balance in patients with peripheral neuropathy. *Arch.Phys.Med.Rehabil.* 82 (2):205-209, 2001.

Level 3:

F. S. Duran, L. Lugo, L. Ramirez, and E. Eusse. Effects of an exercise program on the rehabilitation of patients with spinal cord injury. *Arch.Phys.Med.Rehabil.* 82 (10):1349-1354, 2001.

G. R. Jones, J. M. Jakobi, A. W. Taylor, R. J. Petrella, and A. A. Vandervoort. Community exercise program for older adults recovering from hip fracture: a pilot study. *J.Aging Phys.Act.* 14 (4):439-455, 2006.

J. Jurgel, L. Rannama, H. Gapeyeva, J. Ereline, I. Kolts, and M. Paasuke. Shoulder function in patients with frozen shoulder before and after 4-week rehabilitation. *Medicina (Kaunas.)* 41 (1):30-38, 2005.

Z. Matjacic and A. Zupan. Effects of dynamic balance training during standing and stepping in patients with hereditary sensory motor neuropathy. *Disabil.Rehabil.* 28 (23):1455-1459, 2006.

L. Meiworm, E. Jakob, U. A. Walker, H. H. Peter, and J. Keul. Patients with fibromyalgia benefit from aerobic endurance exercise. *Clin.Rheumatol.* 19 (4):253-257, 2000.

L. Moseley. Combined physiotherapy and education is efficacious for chronic low back pain. *Aust.J.Physiother.* 48 (4):297-302, 2002.

Level 4:

A. L. Betker, A. Desai, C. Nett, N. Kapadia, and T. Szturm. Game-based exercises for dynamic short-sitting balance rehabilitation of people with chronic spinal cord and traumatic brain injuries. *Phys.Ther.* 87 (10):1389-1398, 2007.

- C. L. Blum. Chiropractic and pilates therapy for the treatment of adult scoliosis. *J.Manipulative Physiol Ther.* 25 (4):E3, 2002.
- E. S. Bracher, C. I. Almeida, R. R. Almeida, A. C. Duprat, and C. B. Bracher. A combined approach for the treatment of cervical vertigo. *J.Manipulative Physiol Ther.* 23 (2):96-100, 2000.
- M. P. Callahan, T. Pham, I. Rashbaum, H. Pineda, and N. Greenspan. Cardiopulmonary rehabilitation in a patient with Noonan syndrome. *Arch.Phys.Med.Rehabil.* 81 (2):230-232, 2000.
- C. J. Colloca and B. S. Polkinghorn. Chiropractic management of Ehlers-Danlos syndrome: a report of two cases. *J.Manipulative Physiol Ther.* 26 (7):448-459, 2003.
- S. Corna, A. Nardone, A. Prestinari, M. Galante, M. Grasso, and M. Schieppati. Comparison of Cawthorne-Cooksey exercises and sinusoidal support surface translations to improve balance in patients with unilateral vestibular deficit. *Arch.Phys.Med.Rehabil.* 84 (8):1173-1184, 2003.
- P. R. Dornan. Incontinence--an aggressive approach to treatment: a case series. *J.Sci.Med.Sport* 8 (4):458-462, 2005.
- J. R. Ferrantelli, D. E. Harrison, D. D. Harrison, and D. Stewart. Conservative treatment of a patient with previously unresponsive whiplash-associated disorders using clinical biomechanics of posture rehabilitation methods. *J.Manipulative Physiol Ther.* 28 (3):e1-e8, 2005.
- K. M. Gill-Body, R. A. Popat, S. W. Parker, and D. E. Krebs. Rehabilitation of balance in two patients with cerebellar dysfunction. *Phys.Ther.* 77 (5):534-552, 1997.
- W. M. Glasoe, M. K. Allen, B. F. Awtry, and H. J. Yack. Weight-bearing immobilization and early exercise treatment following a grade II lateral ankle sprain. *J.Orthop.Sports Phys.Ther.* 29 (7):394-399, 1999.
- B. A. Goldberg, R. J. Nowinski, and F. A. Matsen, III. Outcome of nonoperative management of full-thickness rotator cuff tears. *Clin.Orthop.Relat Res.* (382):99-107, 2001.
- R. Grant, G. Jull, and T. Spencer. Active stabilisation training for screen based keyboard operators - a single case study. *Aust.J.Physiother.* 43 (4):235-242, 1997.
- B. L. Greene. Physical therapist management of fluoroquinolone-induced Achilles tendinopathy. *Phys.Ther.* 82 (12):1224-1231, 2002.
- C. M. Gregory, M. G. Bowden, A. Jayaraman, P. Shah, A. Behrman, S. A. Kautz, and K. Vandeborne. Resistance training and locomotor recovery after incomplete spinal cord injury: a case series. *Spinal Cord.* 45 (7):522-530, 2007.
- J. W. Haas, D. E. Harrison, D. D. Harrison, and B. Bymers. Conservative treatment of a patient with syringomyelia using chiropractic biophysics protocols. *J.Manipulative Physiol Ther.* 28 (6):452, 2005.
- C. F. Holmes, J. P. Fletcher, M. J. Blaschak, and R. C. Schenck. Management of shoulder dysfunction with an alternative model of orthopaedic physical therapy intervention: a case report. *J.Orthop.Sports Phys.Ther.* 26 (6):347-354, 1997.

- C. A. Johnston, D. M. Lindsay, and J. P. Wiley. Treatment of iliopsoas syndrome with a hip rotation strengthening program: a retrospective case series. *J.Orthop.Sports Phys.Ther.* 29 (4):218-224, 1999.
- W. B. Katzman, D. E. Sellmeyer, A. L. Stewart, L. Wanek, and K. A. Hamel. Changes in flexed posture, musculoskeletal impairments, and physical performance after group exercise in community-dwelling older women. *Arch.Phys.Med.Rehabil.* 88 (2):192-199, 2007.
- K. Kawanabe, A. Kawashima, I. Sashimoto, T. Takeda, Y. Sato, and J. Iwamoto. Effect of whole-body vibration exercise and muscle strengthening, balance, and walking exercises on walking ability in the elderly. *Keio J.Med.* 56 (1):28-33, 2007.
- P. Kluding and P. Q. McGinnis. Multidimensional exercise for people with Parkinson's disease: a case report. *Physiother.Theory.Pract.* 22 (3):153-162, 2006.
- R. A. Liusuwan, L. M. Widman, R. T. Abresch, A. J. Johnson, and C. M. McDonald. Behavioral intervention, exercise, and nutrition education to improve health and fitness (BENEFit) in adolescents with mobility impairment due to spinal cord dysfunction. *J.Spinal Cord.Med.* 30 Suppl 1:S119-S126, 2007.
- C. W. MacDonald, J. M. Whitman, J. A. Cleland, M. Smith, and H. L. Hoeksma. Clinical outcomes following manual physical therapy and exercise for hip osteoarthritis: A case series. *J.Orthop.Sports Phys.Ther.* 36 (8):588-599, 2006.
- K. K. Mangione and K. M. Palombaro. Exercise prescription for a patient 3 months after hip fracture. *Phys.Ther.* 85 (7):676-687, 2005.
- C. Y. Mao, W. C. Jaw, and H. C. Cheng. Frozen shoulder: correlation between the response to physical therapy and follow-up shoulder arthrography. *Arch.Phys.Med.Rehabil.* 78 (8):857-859, 1997.
- P. W. Marshall and B. A. Murphy. Evaluation of functional and neuromuscular changes after exercise rehabilitation for low back pain using a Swiss ball: a pilot study. *J.Manipulative Physiol Ther.* 29 (7):550-560, 2006.
- M. W. Morningstar, D. Woggon, and G. Lawrence. Scoliosis treatment using a combination of manipulative and rehabilitative therapy: a retrospective case series. *BMC.Musculoskelet.Disord.* 5:32, 2004.
- M. W. Morningstar and T. Joy. Scoliosis treatment using spinal manipulation and the Pettibon Weighting System: a summary of 3 atypical presentations. *Chiropr.Osteopat.* 14:1, 2006.
- M. W. Morningstar. Improvement of lower extremity electrodiagnostic findings following a trial of spinal manipulation and motion-based therapy. *Chiropr.Osteopat.* 14:20, 2006.
- S. Mudge, L. Rochester, and A. Recordon. The effect of treadmill training on gait, balance and trunk control in a hemiplegic subject: a single system design. *Disabil.Rehabil.* 25 (17):1000-1007, 2003.
- G. P. Paulk and D. E. Harrison. Management of a chronic lumbar disk herniation with chiropractic biophysics methods after failed chiropractic manipulative intervention. *J.Manipulative Physiol Ther.* 27 (9):579, 2004.

M. Rahlin. TAMO therapy as a major component of physical therapy intervention for an infant with congenital muscular torticollis: a case report. *Pediatr.Phys.Ther.* 17 (3):209-218, 2005.

J. Rainville, C. Hartigan, C. Jouve, and E. Martinez. The influence of intense exercise-based physical therapy program on back pain anticipated before and induced by physical activities. *Spine J.* 4 (2):176-183, 2004.

S. E. Ross and K. M. Guskiewicz. Effect of coordination training with and without stochastic resonance stimulation on dynamic postural stability of subjects with functional ankle instability and subjects with stable ankles. *Clin.J.Sport Med.* 16 (4):323-328, 2006.

G. Schalow and P. Jaigma. Improvement after severe traumatic brain injury induced by coordination dynamics therapy: comparison with physiologic CNS development. *Electromyogr.Clin.Neurophysiol.* 46 (4):195-209, 2006.

L. Sheppard, H. Mudie, and E. Froude. An investigation of bilateral isokinematic training and neurodevelopmental therapy in improving use of the affected hand in children with hemiplegia. *Phys.Occup.Ther.Pediatr.* 27 (1):5-25, 2007.

D. D. Sherry, C. A. Wallace, C. Kelley, M. Kidder, and L. Sapp. Short- and long-term outcomes of children with complex regional pain syndrome type I treated with exercise therapy. *Clin.J.Pain* 15 (3):218-223, 1999.

B. A. Smith, M. Kubo, D. P. Black, K. G. Holt, and B. D. Ulrich. Effect of practice on a novel task--walking on a treadmill: preadolescents with and without Down syndrome. *Phys.Ther.* 87 (6):766-777, 2007.

T. M. Steffen, B. F. Boeve, L. A. Mollinger-Riemann, and C. M. Petersen. Long-term locomotor training for gait and balance in a patient with mixed progressive supranuclear palsy and corticobasal degeneration. *Phys.Ther.* 87 (8):1078-1087, 2007.

T. A. Torstensen, H. D. Meen, and M. Stiris. The effect of medical exercise therapy on a patient with chronic supraspinatus tendinitis. Diagnostic ultrasound--tissue regeneration: a case study. *J.Orthop.Sports Phys.Ther.* 20 (6):319-327, 1994.

J. Y. Tsauo, H. Y. Lee, J. H. Hsu, C. Y. Chen, and C. J. Chen. Physical exercise and health education for neck and shoulder complaints among sedentary workers. *J.Rehabil.Med.* 36 (6):253-257, 2004.

W. Y. Wang and J. J. Chang. Effects of jumping skill training on walking balance for children with mental retardation and Down's syndrome. *Kaohsiung.J.Med.Sci.* 13 (8):487-495, 1997.

S. L. Whitney, D. M. Wrisley, K. E. Brown, and J. M. Furman. Physical therapy for migraine-related vestibulopathy and vestibular dysfunction with history of migraine. *Laryngoscope* 110 (9):1528-1534, 2000.

S. H. Wiggers and A. Finset. [Rehabilitation of chronic myofascial pain disorders]. *Tidsskr.Nor Laegeforen.* 127 (5):604-608, 2007.

D. Yang, Y. K. Vandongen, and M. C. Stacey. Effect of exercise on calf muscle pump function in patients with chronic venous disease. *Br.J.Surg.* 86 (3):338-341, 1999.

Therapeutic Exercise:

Negative:

Level 1:

S. Crette, H. Moffet, J. Tardif, L. Bessette, F. Morin, P. Fremont, V. Bykerk, C. Thorne, M. Bell, W. Bensen, and C. Blanchette. Intraarticular corticosteroids, supervised physiotherapy, or a combination of the two in the treatment of adhesive capsulitis of the shoulder: a placebo-controlled trial. *Arthritis Rheum.* 48 (3):829-838, 2003.

G. Coughlan and B. Caulfield. A 4-week neuromuscular training program and gait patterns at the ankle joint. *J.Athl.Train.* 42 (1):51-59, 2007.

T. E. Howe, I. Taylor, P. Finn, and H. Jones. Lateral weight transference exercises following acute stroke: a preliminary study of clinical effectiveness. *Clin.Rehabil.* 19 (1):45-53, 2005.

Level 2-3: (0 references)

Level 4:

J. B. Bussmann, M. P. Garssen, P. A. van Doorn, and H. J. Stam. Analysing the favourable effects of physical exercise: relationships between physical fitness, fatigue and functioning in Guillain-Barre syndrome and chronic inflammatory demyelinating polyneuropathy. *J.Rehabil.Med.* 39 (2):121-125, 2007.

Low Level Laser:

Positive:

Level 1:

U. Bingol, L. Altan, and M. Yurtkuran. Low-power laser treatment for shoulder pain. *Photomed.Laser Surg.* 23 (5):459-464, 2005.

J. M. Bjordal, R. A. Lopes-Martins, and V. V. Iversen. A randomised, placebo controlled trial of low level laser therapy for activated Achilles tendinitis with microdialysis measurement of peritendinous prostaglandin E2 concentrations. *Br.J.Sports Med.* 40 (1):76-80, 2006.

R. T. Chow, G. Z. Heller, and L. Barnsley. The effect of 300 mW, 830 nm laser on chronic neck pain: a double-blind, randomized, placebo-controlled study. *Pain* 124 (1-2):201-210, 2006.

G. E. Djavid, R. Mehrdad, M. Ghasemi, H. Hasan-Zadeh, A. Sotoodeh-Manesh, and G. Pouryaghoub. In chronic low back pain, low level laser therapy combined with exercise is more beneficial than exercise alone in the long term: a randomised trial. *Aust.J.Physiother.* 53 (3):155-160, 2007.

- A. Ekim, O. Armagan, F. Tascioglu, C. Oner, and M. Colak. Effect of low level laser therapy in rheumatoid arthritis patients with carpal tunnel syndrome. *Swiss.Med.Wkly.* 137 (23-24):347-352, 2007.
- T. F. Elwakil, A. Elazzazi, and H. Shokeir. Treatment of carpal tunnel syndrome by low-level laser versus open carpal tunnel release. *Lasers Med.Sci.* 22 (4):265-270, 2007.
- D. Evcik, V. Kavuncu, T. Cakir, V. Subasi, and M. Yaman. Laser therapy in the treatment of carpal tunnel syndrome: a randomized controlled trial. *Photomed.Laser Surg.* 25 (1):34-39, 2007.
- S. Fernando, C. M. Hill, and R. Walker. A randomised double blind comparative study of low level laser therapy following surgical extraction of lower third molar teeth. *Br.J.Oral Maxillofac.Surg.* 31 (3):170-172, 1993.
- H. Fikackova, T. Dostalova, L. Navratil, and J. Klaschka. Effectiveness of low-level laser therapy in temporomandibular joint disorders: a placebo-controlled study. *Photomed.Laser Surg.* 25 (4):297-303, 2007.
- A. Gur, M. Karakoc, K. Nas, R. Cevik, J. Sarac, and S. Ataoglu. Effects of low power laser and low dose amitriptyline therapy on clinical symptoms and quality of life in fibromyalgia: a single-blind, placebo-controlled trial. *Rheumatol.Int.* 22 (5):188-193, 2002.
- A. Gur, M. Karakoc, K. Nas, R. Cevik, J. Sarac, and E. Demir. Efficacy of low power laser therapy in fibromyalgia: a single-blind, placebo-controlled trial. *Lasers Med.Sci.* 17 (1):57-61, 2002.
- A. Gur, A. Cosut, A. J. Sarac, R. Cevik, K. Nas, and A. Uyar. Efficacy of different therapy regimes of low-power laser in painful osteoarthritis of the knee: a double-blind and randomized-controlled trial. *Lasers Surg.Med.* 33 (5):330-338, 2003.
- A. Gur, M. Karakoc, R. Cevik, K. Nas, A. J. Sarac, and M. Karakoc. Efficacy of low power laser therapy and exercise on pain and functions in chronic low back pain. *Lasers Surg.Med.* 32 (3):233-238, 2003.
- A. Hakguder, M. Birtane, S. Gurcan, S. Kokino, and F. N. Turan. Efficacy of low level laser therapy in myofascial pain syndrome: an algometric and thermographic evaluation. *Lasers Surg.Med.* 33 (5):339-343, 2003.
- M. Hirschl, R. Katzenschlager, K. Ammer, P. Melnizky, O. Rathkolb, and M. Kundi. Double-blind, randomised, placebo controlled low level laser therapy study in patients with primary Raynaud's phenomenon. *Vasa* 31 (2):91-94, 2002.
- M. Hirschl, R. Katzenschlager, C. Francesconi, and M. Kundi. Low level laser therapy in primary Raynaud's phenomenon--results of a placebo controlled, double blind intervention study. *J.Rheumatol.* 31 (12):2408-2412, 2004.
- J. T. Hopkins, T. A. McLoda, J. G. Seegmiller, and Baxter G. David. Low-Level Laser Therapy Facilitates Superficial Wound Healing in Humans: A Triple-Blind, Sham-Controlled Study. *J.Athl.Train.* 39 (3):223-229, 2004.
- J. Irvine, S. L. Chong, N. Amirjani, and K. M. Chan. Double-blind randomized controlled trial of low-level laser therapy in carpal tunnel syndrome. *Muscle Nerve* 30 (2):182-187, 2004.

M. B. Kreisler, H. A. Haj, N. Noroozi, and B. Willershausen. Efficacy of low level laser therapy in reducing postoperative pain after endodontic surgery-- a randomized double blind clinical study. *Int.J.Oral Maxillofac.Surg.* 33 (1):38-41, 2004.

S. Kulekcioglu, K. Sivrioglu, O. Ozcan, and M. Parlak. Effectiveness of low-level laser therapy in temporomandibular disorder. *Scand.J.Rheumatol.* 32 (2):114-118, 2003.

M. O. Mazzetto, T. G. Carrasco, E. F. Bidinelo, R. C. Andrade Pizzo, and R. G. Mazzetto. Low intensity laser application in temporomandibular disorders: a phase I double-blind study. *Cranio.* 25 (3):186-192, 2007.

M. A. Naeser, K. A. Hahn, B. E. Lieberman, and K. F. Branco. Carpal tunnel syndrome pain treated with low-level laser and microamperes transcutaneous electric nerve stimulation: A controlled study. *Arch.Phys.Med.Rehabil.* 83 (7):978-988, 2002.

O. Oken, Y. Kahraman, F. Ayhan, S. Canpolat, Z. R. Yorgancioglu, and O. F. Oken. The short-term efficacy of laser, brace, and ultrasound treatment in lateral epicondylitis: a prospective, randomized, controlled trial. *J.Hand Ther.* 21 (1):63-67, 2008.

Z. Simunovic, T. Trobonjaca, and Z. Trobonjaca. Treatment of medial and lateral epicondylitis--tennis and golfer's elbow--with low level laser therapy: a multicenter double blind, placebo-controlled clinical study on 324 patients. *J.Clin.Laser Med.Surg.* 16 (3):145-151, 1998.

A. Stergioulas. Low-level laser treatment can reduce edema in second degree ankle sprains. *J.Clin.Laser Med.Surg.* 22 (2):125-128, 2004.

A. Stergioulas. Effects of low-level laser and plyometric exercises in the treatment of lateral epicondylitis. *Photomed.Laser Surg.* 25 (3):205-213, 2007.

A. Stergioulas, M. Stergioula, R. Aarskog, R. A. Lopes-Martins, and J. M. Bjordal. Effects of Low-Level Laser Therapy and Eccentric Exercises in the Treatment of Recreational Athletes With Chronic Achilles Tendinopathy. *Am.J.Sports Med.*, 2008.

Level 2:

M. Milojevic and V. Kuruc. [Low power laser biostimulation in the treatment of bronchial asthma]. *Med.Pregl.* 56 (9-10):413-418, 2003.

Level 3:

Z. Peric. [Influence of low-intensity laser therapy on spatial perception threshold and electroneurographic finding in patients with diabetic polyneuropathy]. *Srp.Ar.h.Celok.Lek.* 135 (5-6):257-263, 2007.

Level 4:

M. Al Awami, M. Schillinger, M. E. Gschwandtner, T. Maca, M. Haumer, and E. Minar. Low level laser treatment of primary and secondary Raynaud's phenomenon. *Vasa* 30 (4):281-284, 2001.

H. Fikackova, T. Dostalova, R. Vosicka, V. Peterova, L. Navratil, and J. Lesak. Arthralgia of the temporomandibular joint and low-level laser therapy. *Photomed.Laser Surg.* 24 (4):522-527, 2006.

S. Giavelli, G. Fava, G. Castronuovo, L. Spinoglio, and A. Galanti. [Low-level laser therapy in osteoarticular diseases in geriatric patients]. *Radiol.Med.(Torino)* 95 (4):303-309, 1998.

P. Monteforte, L. Baratto, L. Molfetta, and G. Rovetta. Low-power laser in osteoarthritis of the cervical spine. *Int.J.Tissue React.* 25 (4):131-136, 2003.

T. Ozen, K. Orhan, I. Gorur, and A. Ozturk. Efficacy of low level laser therapy on neurosensory recovery after injury to the inferior alveolar nerve. *Head Face.Med.* 2:3, 2006.

A. L. Pinheiro, E. T. Cavalcanti, T. I. Pinheiro, M. J. Alves, and C. T. Manzi. Low-level laser therapy in the management of disorders of the maxillofacial region. *J.Clin.Laser Med.Surg.* 15 (4):181-183, 1997.

Z. Simunovic. Low level laser therapy with trigger points technique: a clinical study on 243 patients. *J.Clin.Laser Med.Surg.* 14 (4):163-167, 1996.

Low Level Laser:

Negative:

Level 1:

A. H. Bakhtiary and A. Rashidy-Pour. Ultrasound and laser therapy in the treatment of carpal tunnel syndrome. *Aust.J.Physiother.* 50 (3):147-151, 2004.

L. Brosseau, G. Wells, S. Marchand, I. Gaboury, B. Stokes, M. Morin, L. Casimiro, K. Yonge, and P. Tugwell. Randomized controlled trial on low level laser therapy (LLLT) in the treatment of osteoarthritis (OA) of the hand. *Lasers Surg.Med.* 36 (3):210-219, 2005.

P. C. Conti. Low level laser therapy in the treatment of temporomandibular disorders (TMD): a double-blind pilot study. *Cranio.* 15 (2):144-149, 1997.

R. A. de Bie, H. C. de Vet, T. F. Lenssen, F. A. van den Wildenberg, G. Kootstra, and P. G. Knipschild. Low-level laser therapy in ankle sprains: a randomized clinical trial. *Arch.Phys.Med.Rehabil.* 79 (11):1415-1420, 1998.

J. Hall, A. K. Clarke, D. M. Elvins, and E. F. Ring. Low level laser therapy is ineffective in the management of rheumatoid arthritic finger joints. *Br.J.Rheumatol.* 33 (2):142-147, 1994.

L. A. Matsutani, A. P. Marques, E. A. Ferreira, A. Assumpcao, L. V. Lage, R. A. Casarotto, and C. A. Pereira. Effectiveness of muscle stretching exercises with and without laser therapy at tender points for patients with fibromyalgia. *Clin.Exp.Rheumatol.* 25 (3):410-415, 2007.

D. Mulcahy, D. McCormack, J. McElwain, S. Wagstaff, and C. Conroy. Low level laser therapy: a prospective double blind trial of its use in an orthopaedic population. *Injury* 26 (5):315-317, 1995.

F. Tascioglu, O. Armagan, Y. Tabak, I. Corapci, and C. Oner. Low power laser treatment in patients with knee osteoarthritis. *Swiss.Med.Wkly.* 134 (17-18):254-258, 2004.

H. Thorsen, A. N. Gam, H. Jensen, L. Hojmark, and L. Wahlstrom. [Low energy laser treatment--effect in localized fibromyalgia in the neck and shoulder regions]. *Ugeskr.Laeger* 153 (25):1801-1804, 1991.

H. Thorsen, A. N. Gam, B. H. Svensson, M. Jess, M. K. Jensen, I. Piculell, L. K. Schack, and K. Skjott. Low level laser therapy for myofascial pain in the neck and shoulder girdle. A double-blind, cross-over study. *Scand.J.Rheumatol.* 21 (3):139-141, 1992.

O. Vasseljen, Jr., N. Hoeg, B. Kjeldstad, A. Johnsson, and S. Larsen. Low level laser versus placebo in the treatment of tennis elbow. *Scand.J.Rehabil.Med.* 24 (1):37-42, 1992.

Rde A. Venancio, C. M. Camparis, and Rde F. Lizarelli. Low intensity laser therapy in the treatment of temporomandibular disorders: a double-blind study. *J.Oral Rehabil.* 32 (11):800-807, 2005.

L. H. Zinman, M. Ngo, E. T. Ng, K. T. Nwe, S. Gogov, and V. Bril. Low-intensity laser therapy for painful symptoms of diabetic sensorimotor polyneuropathy: a controlled trial. *Diabetes Care* 27 (4):921-924, 2004.

Level 2-4: (0 references)

TENS:

Positive:

Level 1:

K. Abelson, G. B. Langley, H. Sheppard, M. Vlieg, and R. D. Wigley. Transcutaneous electrical nerve stimulation in rheumatoid arthritis. *N.Z.Med.J.* 96 (727):156-158, 1983.

G. Aydin, S. Tomruk, I. Keles, S. O. Demir, and S. Orkun. Transcutaneous electrical nerve stimulation versus baclofen in spasticity: clinical and electrophysiologic comparison. *Am.J.Phys.Med.Rehabil.* 84 (8):584-592, 2005.

S. Ballegaard, S. J. Christophersen, S. G. Dawids, J. Hesse, and N. V. Olsen. Acupuncture and transcutaneous electric nerve stimulation in the treatment of pain associated with chronic pancreatitis. A randomized study. *Scand.J.Gastroenterol.* 20 (10):1249-1254, 1985.

A. Bertalanffy, A. Kober, P. Bertalanffy, B. Gustorff, O. Gore, S. Adel, and K. Hoerauf. Transcutaneous electrical nerve stimulation reduces acute low back pain during emergency transport. *Acad.Emerg.Med.* 12 (7):607-611, 2005.

E. K. Chee and H. Walton. Treatment of trigger points with microamperage transcutaneous electrical nerve stimulation (TENS)--(the Electro-Acuscope 80). *J.Manipulative Physiol Ther.* 9 (2):131-134, 1986.

G. L. Cheing and C. W. Hui-Chan. Transcutaneous electrical nerve stimulation: nonparallel antinociceptive effects on chronic clinical pain and acute experimental pain. *Arch.Phys.Med.Rehabil.* 80 (3):305-312, 1999.

G. L. Cheing, C. W. Hui-Chan, and K. M. Chan. Does four weeks of TENS and/or isometric exercise produce cumulative reduction of osteoarthritic knee pain? *Clin.Rehabil.* 16 (7):749-760, 2002.

G. L. Cheing and C. W. Hui-Chan. Would the addition of TENS to exercise training produce better physical performance outcomes in people with knee osteoarthritis than either intervention alone? *Clin.Rehabil.* 18 (5):487-497, 2004.

G. L. Cheing and M. L. Luk. Transcutaneous electrical nerve stimulation for neuropathic pain. *J.Hand Surg.[Br.]* 30 (1):50-55, 2005.

T. T. Chiu, C. W. Hui-Chan, and G. Chein. A randomized clinical trial of TENS and exercise for patients with chronic neck pain. *Clin.Rehabil.* 19 (8):850-860, 2005.

M. Y. Dawood and J. Ramos. Transcutaneous electrical nerve stimulation (TENS) for the treatment of primary dysmenorrhea: a randomized crossover comparison with placebo TENS and ibuprofen. *Obstet.Gynecol.* 75 (4):656-660, 1990.

S. Farina, M. Casarotto, M. Benelle, M. Tinazzi, A. Fiaschi, M. Goldoni, and N. Smania. A randomized controlled study on the effect of two different treatments (FREMS AND TENS) in myofascial pain syndrome. *Eura.Medicophys.* 40 (4):293-301, 2004.

V. Finsen, L. Persen, M. Lovlien, E. K. Veslegaard, M. Simensen, A. K. Gasvann, and P. Benum. Transcutaneous electrical nerve stimulation after major amputation. *J.Bone Joint Surg.Br.* 70 (1):109-112, 1988.

T. Forst, M. Nguyen, S. Forst, B. Disselhoff, T. Pohlmann, and A. Pfutzner. Impact of low frequency transcutaneous electrical nerve stimulation on symptomatic diabetic neuropathy using the new Salutaris device. *Diabetes Nutr.Metab* 17 (3):163-168, 2004.

S. B. Graff-Radford, J. L. Reeves, R. L. Baker, and D. Chiu. Effects of transcutaneous electrical nerve stimulation on myofascial pain and trigger point sensitivity. *Pain* 37 (1):1-5, 1989.

D. J. Grant, J. Bishop-Miller, D. M. Winchester, M. Anderson, and S. Faulkner. A randomized comparative trial of acupuncture versus transcutaneous electrical nerve stimulation for chronic back pain in the elderly. *Pain* 82 (1):9-13, 1999.

P. Hansson and A. Eklom. Transcutaneous electrical nerve stimulation (TENS) as compared to placebo TENS for the relief of acute oro-facial pain. *Pain* 15 (2):157-165, 1983.

J. Katz, C. France, and R. Melzack. An association between phantom limb sensations and stump skin conductance during transcutaneous electrical nerve stimulation (TENS) applied to the contralateral leg: a case study. *Pain* 36 (3):367-377, 1989.

A. J. Koke, J. S. Schouten, M. J. Lamerichs-Geelen, J. S. Lipsch, E. M. Waltje, M. van Kleef, and J. Patijn. Pain reducing effect of three types of transcutaneous electrical nerve stimulation in patients with chronic pain: a randomized crossover trial. *Pain* 108 (1-2):36-42, 2004.

T. Lang, R. Barker, B. Steinlechner, B. Gustorff, T. Puskas, O. Gore, and A. Kober. TENS relieves acute posttraumatic hip pain during emergency transport. *J.Trauma* 62 (1):184-188, 2007.

T. R. Lehmann, D. W. Russell, K. F. Spratt, H. Colby, Y. K. Liu, M. L. Fairchild, and S. Christensen. Efficacy of electroacupuncture and TENS in the rehabilitation of chronic low back pain patients. *Pain* 26 (3):277-290, 1986.

T. Lundeberg. The pain suppressive effect of vibratory stimulation and transcutaneous electrical nerve stimulation (TENS) as compared to aspirin. *Brain Res.* 294 (2):201-209, 1984.

C. Mannheimer, S. Lund, and C. A. Carlsson. The effect of transcutaneous electrical nerve stimulation (TNS) on joint pain in patients with rheumatoid arthritis. *Scand.J.Rheumatol.* 7 (1):13-16, 1978.

S. Marchand, J. Charest, J. Li, J. R. Chenard, B. Lavignolle, and L. Laurencelle. Is TENS purely a placebo effect? A controlled study on chronic low back pain. *Pain* 54 (1):99-106, 1993.

R. Melzack, P. Vetere, and L. Finch. Transcutaneous electrical nerve stimulation for low back pain. A comparison of TENS and massage for pain and range of motion. *Phys.Ther.* 63 (4):489-493, 1983.

S. R. Moore and J. Shurman. Combined neuromuscular electrical stimulation and transcutaneous electrical nerve stimulation for treatment of chronic back pain: a double-blind, repeated measures comparison. *Arch.Phys.Med.Rehabil.* 78 (1):55-60, 1997.

Nunez C. Navarro and Carrasco M. Pacheco. [Transcutaneous electric stimulation (TENS) to reduce pain after cesarean section]. *Ginecol.Obstet.Mex.* 68:60-63, 2000.

S. S. Ng and C. W. Hui-Chan. Transcutaneous electrical nerve stimulation combined with task-related training improves lower limb functions in subjects with chronic stroke. *Stroke* 38 (11):2953-2959, 2007.

G. J. Ordog. Transcutaneous electrical nerve stimulation versus oral analgesic: a randomized double-blind controlled study in acute traumatic pain. *Am.J.Emerg.Med.* 5 (1):6-10, 1987.

N. Paker, D. Tekdos, N. Kesiktas, and D. Soy. Comparison of the therapeutic efficacy of TENS versus intra-articular hyaluronic acid injection in patients with knee osteoarthritis: a prospective randomized study. *Adv.Ther.* 23 (2):342-353, 2006.

T. Rahko and V. Kotti. Tinnitus treatment by transcutaneous nerve stimulation (TNS). *Acta Otolaryngol.Suppl* 529:88-89, 1997.

J. P. Sloan, C. L. Muwanga, E. A. Waters, A. F. Dove, and S. H. Dave. Multiple rib fractures: transcutaneous nerve stimulation versus conventional analgesia. *J.Trauma* 26 (12):1120-1122, 1986.

C. M. Smith, M. S. Guralnick, M. M. Gelfand, and M. E. Jeans. The effects of transcutaneous electrical nerve stimulation on post-cesarean pain. *Pain* 27 (2):181-193, 1986.

J. O. Sodipo, S. A. Adedeji, and O. Olumide. Postoperative pain relief by transcutaneous electrical nerve stimulation (TENS). *Am.J.Chin Med.* 8 (1-2):190-194, 1980.

Y. Tekeoglu, B. Adak, and T. Goksoy. Effect of transcutaneous electrical nerve stimulation (TENS) on Barthel Activities of Daily Living (ADL) index score following stroke. *Clin.Rehabil.* 12 (4):277-280, 1998.

M. Tinazzi, S. Farina, K. Bhatia, A. Fiaschi, G. Moretto, L. Bertolasi, S. Zarattini, and N. Smania. TENS for the treatment of writer's cramp dystonia: a randomized, placebo-controlled study. *Neurology* 64 (11):1946-1948, 2005.

H. Tsukayama, H. Yamashita, H. Amagai, and Y. Tanno. Randomised controlled trial comparing the effectiveness of electroacupuncture and TENS for low back pain: a preliminary study for a pragmatic trial. *Acupunct.Med.* 20 (4):175-180, 2002.

N. Tugay, T. Akbayrak, F. Demirturk, I. C. Karakaya, O. Kocaacar, U. Tugay, M. G. Karakaya, and F. Demirturk. Effectiveness of transcutaneous electrical nerve stimulation and interferential current in primary dysmenorrhea. *Pain Med.* 8 (4):295-300, 2007.

C. A. Warfield, J. M. Stein, and H. A. Frank. The effect of transcutaneous electrical nerve stimulation on pain after thoracotomy. *Ann.Thorac.Surg.* 39 (5):462-465, 1985.

K. S. Xu, L. He, J. L. Li, and J. N. Mai. [Effects of transcutaneous electrical nerve stimulation on motor function in ambulant children with spastic cerebral palsy: a randomized trial]. *Zhonghua Er.Ke.Za Zhi.* 45 (8):564-567, 2007.

M. Yurtkuran and T. Kocagil. TENS, electroacupuncture and ice massage: comparison of treatment for osteoarthritis of the knee. *Am.J.Acupunct.* 27 (3-4):133-140, 1999.

Level 2:

C. Herraiz, A. Toledano, and I. Diges. Trans-electrical nerve stimulation (TENS) for somatic tinnitus. *Prog.Brain Res.* 166:389-394, 2007.

Level 3:

J. T. van der Spank, D. C. Cambier, H. M. De Paepe, L. A. Danneels, E. E. Witvrouw, and L. Beerens. Pain relief in labour by transcutaneous electrical nerve stimulation (TENS). *Arch.Gynecol.Obstet.* 264 (3):131-136, 2000.

Level 4:

D. Avdic and A. Buljina. [TENS in the treatment of muscle spasm]. *Med.Arh.* 54 (1):49-51, 2000.

G. Aydemir, M. S. Tezer, P. Borman, H. Bodur, and A. Unal. Treatment of tinnitus with transcutaneous electrical nerve stimulation improves patients' quality of life. *J.Laryngol.Otol.* 120 (6):442-445, 2006.

L. Blumenfeld, Y. Hahn, A. Lepage, R. Leonard, and P. C. Belafsky. Transcutaneous electrical stimulation versus traditional dysphagia therapy: a nonconcurrent cohort study. *Otolaryngol.Head Neck Surg.* 135 (5):754-757, 2006.

E. Bohm. Transcutaneous electrical nerve stimulation in chronic pain after peripheral nerve injury. *Acta Neurochir.(Wien.)* 40 (3-4):277-283, 1978.

A. Bremerich, W. Wiegel, T. Thein, and T. Dietze. Transcutaneous electric nerve stimulation (TENS) in the therapy of chronic facial pain. Preliminary report. *J.Craniomaxillofac.Surg.* 16 (8):379-381, 1988.

R. A. Carabelli and W. C. Kellerman. Phantom limb pain: relief by application of TENS to contralateral extremity. *Arch.Phys.Med.Rehabil.* 66 (7):466-467, 1985.

B. Casa, R. Galvagni, R. Marsigli, A. Ape, G. Manzoni, and A. Panicalli. [Usefulness of transcutaneous electric nerve stimulation in the treatment of headache of cervical origin]. *Acta Biomed.Ateneo.Parmense.* 57 (3-4):89-108, 1986.

R. Davis and R. Lentini. Transcutaneous nerve stimulation for treatment of pain in patients with spinal cord injury. *Surg.Neurol.* 4 (1):100-101, 1975.

A. S. Dil'din, L. I. Kalakutskii, and S. L. Nesterov. [Transcutaneous electrostimulation and its place in the treatment of chronic pain syndromes]. *Zh.Nevropatol.Psikhiatr.Im S.S.Korsakova* 84 (4):539-542, 1984.

T. A. Fredriksen, S. Bergmann, J. P. Hesselberg, A. Stolt-Nielsen, R. Ringkjøb, and O. Sjaastad. Electrical stimulation in multiple sclerosis. Comparison of transcutaneous electrical stimulation and epidural spinal cord stimulation. *Appl.Neurophysiol.* 49 (1-2):4-24, 1986.

J. L. Hollinger. Transcutaneous electrical nerve stimulation after cesarean birth. *Phys.Ther.* 66 (1):36-38, 1986.

F. Jacobsson, A. Himmelmann, A. Bergbrant, A. Svensson, and C. Mannheimer. The effect of transcutaneous electric nerve stimulation in patients with therapy-resistant hypertension. *J.Hum.Hypertens.* 14 (12):795-798, 2000.

B. Kaada. Treatment of peritendinitis calcarea of the shoulder by transcutaneous nerve stimulation. *Acupunct.Electrother.Res.* 9 (2):115-125, 1984.

B. Kaada. Systemic sclerosis: successful treatment of ulcerations, pain, Raynaud's phenomenon, calcinosis, and dysphagia by transcutaneous nerve stimulation. A case report. *Acupunct.Electrother.Res.* 9 (1):31-44, 1984.

B. Kaada, S. Hognestad, and J. Havstad. Transcutaneous nerve stimulation (TNS) in tinnitus. *Scand.Audiol.* 18 (4):211-217, 1989.

J. Kahn. Transcutaneous electrical nerve stimulation for nonunited fractures; a clinical report. *Phys.Ther.* 62 (6):840-844, 1982.

D. J. Krauss and O. M. Lilien. Transcutaneous electrical nerve stimulator for stress incontinence. *J.Urol.* 125 (6):790-793, 1981.

T. Lundeberg. A comparative study of the pain alleviating effect of vibratory stimulation, transcutaneous electrical nerve stimulation, electroacupuncture and placebo. *Am.J.Chin Med.* 12 (1-4):72-79, 1984.

G. J. Magarian, B. Leikam, and R. Palac. Transcutaneous electrical nerve stimulation (TENS) for treatment of severe angina pectoris refractory to maximal medical and surgical management--a case report. *Angiology* 41 (5):408-411, 1990.

E. Malm-Buatsi, K. G. Nepple, M. A. Boyt, J. C. Austin, and C. S. Cooper. Efficacy of transcutaneous electrical nerve stimulation in children with overactive bladder refractory to pharmacotherapy. *Urology* 70 (5):980-983, 2007.

S. Mangold, T. Keller, A. Curt, and V. Dietz. Transcutaneous functional electrical stimulation for grasping in subjects with cervical spinal cord injury. *Spinal Cord*. 43 (1):1-13, 2005.

C. Mannheimer, C. A. Carlsson, H. Emanuelsson, A. Vedin, F. Waagstein, and C. Wilhelmsson. The effects of transcutaneous electrical nerve stimulation in patients with severe angina pectoris. *Circulation* 71 (2):308-316, 1985.

C. Mannheimer, C. A. Carlsson, A. Vedin, and C. Wilhelmsson. Transcutaneous electrical nerve stimulation (TENS) in angina pectoris. *Pain* 26 (3):291-300, 1986.

J. A. McCarthy and R. W. Zigenfus. Transcutaneous electrical nerve stimulation: an adjunct in the pain management of Guillain-Barre syndrome. *Phys.Ther.* 58 (1):23-24, 1978.

W. J. Meyler, M. J. de Jongste, and C. A. Rolf. Clinical evaluation of pain treatment with electrostimulation: a study on TENS in patients with different pain syndromes. *Clin.J.Pain* 10 (1):22-27, 1994.

S. Pohl, T. Masyk-Iversen, U. Lips, and I. Pichlmayr. [Transcutaneous electrical nerve stimulation (TENS) in the treatment of a peripheral lesion of the brachial plexus]. *Anasth.Intensivther.Notfallmed.* 18 (3):144-146, 1983.

K. P. Potisk, M. Gregoric, and L. Vodovnik. Effects of transcutaneous electrical nerve stimulation (TENS) on spasticity in patients with hemiplegia. *Scand.J.Rehabil.Med.* 27 (3):169-174, 1995.

F. J. Robaina, J. L. Rodriguez, J. A. de Vera, and M. A. Martin. Transcutaneous electrical nerve stimulation and spinal cord stimulation for pain relief in reflex sympathetic dystrophy. *Stereotact.Funct.Neurosurg.* 52 (1):53-62, 1989.

D. Rodrigues, A. O. Siriani, and F. Berzin. Effect of conventional TENS on pain and electromyographic activity of masticatory muscles in TMD patients. *Braz.Oral Res.* 18 (4):290-295, 2004.

E. Savk, O. Savk, and F. Sendur. Transcutaneous electrical nerve stimulation offers partial relief in notalgia paresthetica patients with a relevant spinal pathology. *J.Dermatol.* 34 (5):315-319, 2007.

G. D. Schuster and M. C. Infante. Pain relief after low back surgery: the efficacy of transcutaneous electrical nerve stimulation. *Pain* 8 (3):299-302, 1980.

D. L. Somers and M. F. Somers. Treatment of neuropathic pain in a patient with diabetic neuropathy using transcutaneous electrical nerve stimulation applied to the skin of the lumbar region. *Phys.Ther.* 79 (8):767-775, 1999.

P. Taylor, M. Hallett, and L. Flaherty. Treatment of osteoarthritis of the knee with transcutaneous electrical nerve stimulation. *Pain* 11 (2):233-240, 1981.

J. U. Toglia and K. Izzo. Treatment of myoclonic dystonia with transcutaneous electrical nerve stimulation. *Ital.J.Neurol.Sci.* 6 (1):75-78, 1985.

TENS:

Negative:

Level 1:

R. J. Cuschieri, C. G. Morran, and C. S. McArdle. Transcutaneous electrical stimulation for postoperative pain. *Ann.R.Coll.Surg.Engl.* 67 (2):127-129, 1985.

R. A. Deyo, N. E. Walsh, D. C. Martin, L. S. Schoenfeld, and S. Ramamurthy. A controlled trial of transcutaneous electrical nerve stimulation (TENS) and exercise for chronic low back pain. *N.Engl.J.Med.* 322 (23):1627-1634, 1990.

J. M. Gilbert, T. Gledhill, N. Law, and C. George. Controlled trial of transcutaneous electrical nerve stimulation (TENS) for postoperative pain relief following inguinal herniorrhaphy. *Br.J.Surg.* 73 (9):749-751, 1986.

E. Herman, R. Williams, P. Stratford, A. Fargas-Babjak, and M. Trott. A randomized controlled trial of transcutaneous electrical nerve stimulation (CODETRON) to determine its benefits in a rehabilitation program for acute occupational low back pain. *Spine* 19 (5):561-568, 1994.

N. D. Kofotolis, S. P. Vlachopoulos, and E. Kellis. Sequentially allocated clinical trial of rhythmic stabilization exercises and TENS in women with chronic low back pain. *Clin.Rehabil.* 22 (2):99-111, 2008.

P. P. Law, G. L. Cheing, and A. Y. Tsui. Does Transcutaneous Electrical Nerve Stimulation Improve the Physical Performance of People With Knee Osteoarthritis? *J.Clin.Rheumatol.* 10 (6):295-299, 2004.

L. Miller, P. Mattison, L. Paul, and L. Wood. The effects of transcutaneous electrical nerve stimulation (TENS) on spasticity in multiple sclerosis. *Mult.Scler.* 13 (4):527-533, 2007.

E. Sherry, P. Kitchener, and R. Smart. A prospective randomized controlled study of VAX-D and TENS for the treatment of chronic low back pain. *Neurol.Res.* 23 (7):780-784, 2001.

J. F. Stubbing and J. A. Jellicoe. Transcutaneous electrical nerve stimulation after thoracotomy. Pain relief and peak expiratory flow rate--a trial of transcutaneous electrical nerve stimulation. *Anaesthesia* 43 (4):296-298, 1988.

J. M. van der Ploeg, H. A. Vervest, A. L. Liem, and J. H. Schagen van Leeuwen. Transcutaneous nerve stimulation (TENS) during the first stage of labour: a randomized clinical trial. *Pain* 68 (1):75-78, 1996.

K. Warke, J. Al Smadi, D. Baxter, D. M. Walsh, and A. S. Lowe-Strong. Efficacy of transcutaneous electrical nerve stimulation (tens) for chronic low-back pain in a multiple sclerosis population: a randomized, placebo-controlled clinical trial. *Clin.J.Pain* 22 (9):812-819, 2006.

K. Wieselmann-Penkner, M. Janda, M. Lorenzoni, and R. Polansky. A comparison of the muscular relaxation effect of TENS and EMG-biofeedback in patients with bruxism. *J.Oral Rehabil.* 28 (9):849-853, 2001.

Level 2-3: (0 references)

Level 4:

J. J. Choi, K. Srikantha, and W. H. Wu. A comparison of electro-acupuncture, transcutaneous electrical nerve stimulation and laser photo-biostimulation on pain relief and glucocorticoid excretion. A case report. *Acupunct.Electrother.Res.* 11 (1):45-51, 1986.

Whole Body Vibration

Positive:

Level 1:

L. Ahlborg, C. Andersson, and P. Julin. Whole-body vibration training compared with resistance training: effect on spasticity, muscle strength and motor performance in adults with cerebral palsy. *J.Rehabil.Med.* 38 (5):302-308, 2006.

D. L. Belavy, J. A. Hides, S. J. Wilson, W. Stanton, F. C. Dimeo, J. Rittweger, D. Felsenberg, and C. A. Richardson. Resistive simulated weightbearing exercise with whole body vibration reduces lumbar spine deconditioning in bed-rest. *Spine* 33 (5):E121-E131, 2008.

M. Cardinale, J. Leiper, P. Farajian, and M. Heer. Whole-body vibration can reduce calciuria induced by high protein intakes and may counteract bone resorption: A preliminary study. *J.Sports Sci.* 25 (1):111-119, 2007.

C. T. Haas, S. Turbanski, K. Kessler, and D. Schmidtbleicher. The effects of random whole-body-vibration on motor symptoms in Parkinson's disease. *NeuroRehabilitation.* 21 (1):29-36, 2006.

J. Iwamoto, T. Takeda, Y. Sato, and M. Uzawa. Effect of whole-body vibration exercise on lumbar bone mineral density, bone turnover, and chronic back pain in post-menopausal osteoporotic women treated with alendronate. *Aging Clin.Exp.Res.* 17 (2):157-163, 2005.

A. Moezy, G. Olyaei, M. Hadian, M. Razi, and S. Faghihzadeh. A Comparative Study of Whole Body Vibration Training and Conventional Training on Knee Proprioception and Postural Stability after Anterior Cruciate Ligament Reconstruction. *Br.J.Sports Med.*, 2008.

J. Rittweger, K. Just, K. Kautzsch, P. Reeg, and D. Felsenberg. Treatment of chronic lower back pain with lumbar extension and whole-body vibration exercise: a randomized controlled trial. *Spine* 27 (17):1829-1834, 2002.

O. Schuhfried, C. Mittermaier, T. Jovanovic, K. Pieber, and T. Paternostro-Sluga. Effects of whole-body vibration in patients with multiple sclerosis: a pilot study. *Clin.Rehabil.* 19 (8):834-842, 2005.

T. K. Tihanyi, M. Horvath, G. Fazekas, T. Hortobagyi, and J. Tihanyi. One session of whole body vibration increases voluntary muscle strength transiently in patients with stroke. *Clin.Rehabil.* 21 (9):782-793, 2007.

S. M. Verschueren, M. Roelants, C. Delecluse, S. Swinnen, D. Vanderschueren, and S. Boonen. Effect of 6-month whole body vibration training on hip density, muscle strength, and postural control in postmenopausal women: a randomized controlled pilot study. *J.Bone Miner.Res.* 19 (3):352-359, 2004.

Level 2: (0 references)

Level 3:

V. Gilsanz, T. A. Wren, M. Sanchez, F. Dorey, S. Judex, and C. Rubin. Low-level, high-frequency mechanical signals enhance musculoskeletal development of young women with low BMD. *J.Bone Miner.Res.* 21 (9):1464-1474, 2006.

S. Turbanski, C. T. Haas, D. Schmidtbleicher, A. Friedrich, and P. Duisberg. Effects of random whole-body vibration on postural control in Parkinson's disease. *Res.Sports Med.* 13 (3):243-256, 2005.

Level 4:

O. Semler, O. Fricke, K. Vezyroglou, C. Stark, and E. Schoenau. Preliminary results on the mobility after whole body vibration in immobilized children and adolescents. *J.Musculoskelet.Neuronal.Interact.* 7 (1):77-81, 2007.

Whole Body Vibration:

Negative:

Level 1-3: (0 References)

Level 4:

O. Semler, O. Fricke, K. Vezyroglou, C. Stark, and E. Schoenau. Preliminary results on the mobility after whole body vibration in immobilized children and adolescents. *J.Musculoskelet.Neuronal.Interact.* 7 (1):77-81, 2007.

Yoga:

Positive:

Level 1:

G. D. da Silva, G. Lorenzi-Filho, and L. V. Lage. Effects of yoga and the addition of Tui Na in patients with fibromyalgia. *J.Altern.Complement Med.* 13 (10):1107-1113, 2007.

T. Fluge, J. Richter, H. Fabel, E. Zysno, E. Weller, and T. O. Wagner. [Long-term effects of breathing exercises and yoga in patients with bronchial asthma]. *Pneumologie* 48 (7):484-490, 1994.

M. L. Galantino, T. M. Bzdewka, J. L. Eissler-Russo, M. L. Holbrook, E. P. Mogck, P. Geigle, and J. T. Farrar. The impact of modified Hatha yoga on chronic low back pain: a pilot study. *Altern.Ther.Health Med.* 10 (2):56-59, 2004.

M. S. Garfinkel, A. Singhal, W. A. Katz, D. A. Allan, R. Reshetar, and H. R. Schumacher, Jr. Yoga-based intervention for carpal tunnel syndrome: a randomized trial. *JAMA* 280 (18):1601-1603, 1998.

P. J. John, N. Sharma, C. M. Sharma, and A. Kankane. Effectiveness of yoga therapy in the treatment of migraine without aura: a randomized controlled trial. *Headache* 47 (5):654-661, 2007.

L. Kuttner, C. T. Chambers, J. Hardial, D. M. Israel, K. Jacobson, and K. Evans. A randomized trial of yoga for adolescents with irritable bowel syndrome. *Pain Res.Manag.* 11 (4):217-223, 2006.

R. Manocha, G. B. Marks, P. Kenchington, D. Peters, and C. M. Salome. Sahaja yoga in the management of moderate to severe asthma: a randomised controlled trial. *Thorax* 57 (2):110-115, 2002.

R. McCaffrey, P. Ruknui, U. Hatthakit, and P. Kasetsoomboon. The effects of yoga on hypertensive persons in Thailand. *Holist.Nurs.Pract.* 19 (4):173-180, 2005.

A. B. Moadel, C. Shah, J. Wylie-Rosett, M. S. Harris, S. R. Patel, C. B. Hall, and J. A. Sparano. Randomized controlled trial of yoga among a multiethnic sample of breast cancer patients: effects on quality of life. *J.Clin.Oncol.* 25 (28):4387-4395, 2007.

R. Nagarathna and H. R. Nagendra. Yoga for bronchial asthma: a controlled study. *Br.Med.J.(Clin.Res.Ed)* 291 (6502):1077-1079, 1985.

B. S. Oken, S. Kishiyama, D. Zajdel, D. Bourdette, J. Carlsen, M. Haas, C. Hugos, D. F. Kraemer, J. Lawrence, and M. Mass. Randomized controlled trial of yoga and exercise in multiple sclerosis. *Neurology* 62 (11):2058-2064, 2004.

S. Sareen, V. Kumari, K. S. Gajebasia, and N. K. Gajebasia. Yoga: a tool for improving the quality of life in chronic pancreatitis. *World J.Gastroenterol.* 13 (3):391-397, 2007.

D. S. Shannahoff-Khalsa, L. E. Ray, S. Levine, C. C. Gallen, B. J. Schwartz, and J. J. Sidorowich. Randomized controlled trial of yogic meditation techniques for patients with obsessive-compulsive disorder. *CNS.Spectr.* 4 (12):34-47, 1999.

K. J. Sherman, D. C. Cherkin, J. Erro, D. L. Miglioretti, and R. A. Deyo. Comparing yoga, exercise, and a self-care book for chronic low back pain: a randomized, controlled trial. *Ann.Intern.Med.* 143 (12):849-856, 2005.

C. Smith, H. Hancock, J. Blake-Mortimer, and K. Eckert. A randomised comparative trial of yoga and relaxation to reduce stress and anxiety. *Complement Ther.Med.* 15 (2):77-83, 2007.

P. K. Vedanthan, L. N. Kesavalu, K. C. Murthy, K. Duvall, M. J. Hall, S. Baker, and S. Nagarathna. Clinical study of yoga techniques in university students with asthma: a controlled study. *Allergy Asthma Proc.* 19 (1):3-9, 1998.

N. K. Visweswaraiah and S. Telles. Randomized trial of yoga as a complementary therapy for pulmonary tuberculosis. *Respirology.* 9 (1):96-101, 2004.

K. A. Williams, J. Petronis, D. Smith, D. Goodrich, J. Wu, N. Ravi, E. J. Doyle, Jr., Juckett R. Gregory, Kolar M. Munoz, R. Gross, and L. Steinberg. Effect of Iyengar yoga therapy for chronic low back pain. *Pain* 115 (1-2):107-117, 2005.

M. Yurtkuran, A. Alp, M. Yurtkuran, and K. Dilek. A modified yoga-based exercise program in hemodialysis patients: a randomized controlled study. *Complement Ther.Med.* 15 (3):164-171, 2007.

Level 2:

H. R. Nagendra and R. Nagarathna. An integrated approach of yoga therapy for bronchial asthma: a 3-54-month prospective study. *J.Asthma* 23 (3):123-137, 1986.

Level 3:

V. Malhotra, S. Singh, O. P. Tandon, S. V. Madhu, A. Prasad, and S. B. Sharma. Effect of Yoga asanas on nerve conduction in type 2 diabetes. *Indian J.Physiol Pharmacol.* 46 (3):298-306, 2002.

S. Narendran, R. Nagarathna, S. Gunasheela, and H. R. Nagendra. Efficacy of yoga in pregnant women with abnormal Doppler study of umbilical and uterine arteries. *J.Indian Med.Assoc.* 103 (1):12-17, 2005.

Level 4:

J. V. Bastille and K. M. Gill-Body. A yoga-based exercise program for people with chronic poststroke hemiparesis. *Phys.Ther.* 84 (1):33-48, 2004.

D. Behera. Yoga therapy in chronic bronchitis. *J.Assoc.Physicians India* 46 (2):207-208, 1998.

C. Booth-LaForce, R. C. Thurston, and M. R. Taylor. A pilot study of a Hatha yoga treatment for menopausal symptoms. *Maturitas* 57 (3):286-295, 2007.

E. L. Bukowski, A. Conway, L. A. Glentz, K. Kurland, and M. L. Galantino. The effect of iyengar yoga and strengthening exercises for people living with osteoarthritis of the knee: a case series. *Int.Q.Community Health Educ.* 26 (3):287-305, 2006.

S. C. Jain, L. Rai, A. Valecha, U. K. Jha, S. O. Bhatnagar, and K. Ram. Effect of yoga training on exercise tolerance in adolescents with childhood asthma. *J.Asthma* 28 (6):437-442, 1991.

S. L. Kolasinski, M. Garfinkel, A. G. Tsai, W. Matz, A. Van Dyke, and H. R. Schumacher. Iyengar yoga for treating symptoms of osteoarthritis of the knees: a pilot study. *J.Altern.Complement Med.* 11 (4):689-693, 2005.

D. Shapiro, I. A. Cook, D. M. Davydov, C. Ottaviani, A. F. Leuchter, and M. Abrams. Yoga as a Complementary Treatment of Depression: Effects of Traits and Moods on Treatment Outcome. *Evid.Based.Complement Alternat.Med.* 4 (4):493-502, 2007.

S. Singh, V. Malhotra, K. P. Singh, S. V. Madhu, and O. P. Tandon. Role of yoga in modifying certain cardiovascular functions in type 2 diabetic patients. *J.Assoc.Physicians India* 52:203-206, 2004.

S. Sundar, S. K. Agrawal, V. P. Singh, S. K. Bhattacharya, K. N. Udupa, and S. K. Vaish. Role of yoga in management of essential hypertension. *Acta Cardiol.* 39 (3):203-208, 1984.

Yoga:

Negative:

Level 1:

A. B. Sabina, A. L. Williams, H. K. Wall, S. Bansal, G. Chupp, and D. L. Katz. Yoga intervention for adults with mild-to-moderate asthma: a pilot study. *Ann.Allergy Asthma Immunol.* 94 (5):543-548, 2005.

G. A. van Montfrans, J. M. Karemaker, W. Wieling, and A. J. Dunning. Relaxation therapy and continuous ambulatory blood pressure in mild hypertension: a controlled study. *BMJ* 300 (6736):1368-1372, 1990.

Level 2-4: (0 References)

Extension Traction:

Positive:

Level 1:

D. D. Harrison, B. L. Jackson, S. Troyanovich, G. Robertson, D. de George, and W. F. Barker. The efficacy of cervical extension-compression traction combined with diversified manipulation and drop

table adjustments in the rehabilitation of cervical lordosis: a pilot study. *J.Manipulative Physiol Ther.* 17 (7):454-464, 1994.

Level 2:

D. E. Harrison, R. Cailliet, D. D. Harrison, T. J. Janik, and B. Holland. A new 3-point bending traction method for restoring cervical lordosis and cervical manipulation: a nonrandomized clinical controlled trial. *Arch.Phys.Med.Rehabil.* 83 (4):447-453, 2002.

D. E. Harrison, R. Cailliet, D. D. Harrison, T. J. Janik, and B. Holland. Changes in sagittal lumbar configuration with a new method of extension traction: nonrandomized clinical controlled trial. *Arch.Phys.Med.Rehabil.* 83 (11):1585-1591, 2002.

D. E. Harrison, D. D. Harrison, J. J. Betz, T. J. Janik, B. Holland, C. J. Colloca, and J. W. Haas. Increasing the cervical lordosis with chiropractic biophysics seated combined extension-compression and transverse load cervical traction with cervical manipulation: nonrandomized clinical control trial. *J.Manipulative Physiol Ther.* 26 (3):139-151, 2003.

Level 3: (0 references)

Level 4:

J. R. Ferrantelli, D. E. Harrison, D. D. Harrison, and D. Stewart. Conservative treatment of a patient with previously unresponsive whiplash-associated disorders using clinical biomechanics of posture rehabilitation methods. *J.Manipulative Physiol Ther.* 28 (3):e1-e8, 2005.

J. W. Haas, D. E. Harrison, D. D. Harrison, and B. Bymers. Conservative treatment of a patient with syringomyelia using chiropractic biophysics protocols. *J.Manipulative Physiol Ther.* 28 (6):452, 2005.

G. P. Paulk and D. E. Harrison. Management of a chronic lumbar disk herniation with chiropractic biophysics methods after failed chiropractic manipulative intervention. *J.Manipulative Physiol Ther.* 27 (9):579, 2004.

Extension Traction:

Negative:

Level 1-4: (0 References)

References

1. Christensen K. Some Chiropractic Mileposts on the Physiotherapy Timeline. *Dynamic Chiropractic* 15[03]. 1997. Ref Type: Magazine Article
2. Jaskoviak P, Schafer R: *Applied Physiotherapy*, American Chiropractic Association, 1993
3. Massey BF, Jr.: Making vision 2020 a reality. *Phys Ther* 83:1023-1026, 2003
4. Moffat M: The History of Physical Therapy Practice in the United States. *Journal of Physical Therapy Education* 2003

5. Peterson D, Wiese G: Chiropractic: An Illustrated History, 1985
6. Snook G: The history of sports medicine. Part 1. American Journal of Sports Medicine 12:252-254, 1984